

1 TO: Don Spagnolo, General Manager, Nipomo Community Services District
2 FROM: Joel Degner E.I.T., Brad Newton, Ph.D., P.G.
3 RE: Spring 2010 Groundwater Index
4 DATE: July 15, 2010

5 INTRODUCTION

6 Groundwater surface elevations (GSE) underlying the Nipomo Mesa are regularly
7 measured at many places (wells) across the mesa. The Spring 2010 Groundwater Index (GWI)
8 has been computed and presented herein along with historical GWI from 1975 to present based
9 on these groundwater surface elevation measurements collected during spring and fall across
10 the Nipomo Mesa. Limited measurements of GSE were available for the years 1982, 1983, 1984,
11 1994 and 1997, thus precluding a reliable calculation of GWI for those years.

12 Ground elevation surveys for the key wells were conducted in preparation of the 1st
13 Annual Report - Calendar Year 2008 for the Nipomo Mesa Management Area (NMMA). These
14 updated reference points were not incorporated into the GWI to preserve consistency in the
15 historical calculations and presentations.

16 The NMMA Technical Group has not reviewed this technical memorandum, its findings,
17 or any presentation of this evaluation.

18

19 RESULTS

20 Spring 2010 GWI is 80,000 acre-feet (AF), which is 4,000 AF greater than the Spring 2009
21 GWI (Table 1, Figure 1). The Key Well Index from NMMA 2nd Annual Report - Calendar Year
22 2009 generally follows the same historical trends as the GWI (Figure 1).

23

24 METHODOLOGY

25 The calculation of Spring and Fall GWI are based on GSE measurements regularly made
26 by San Luis Obispo County Department of Public Works (SLO DPW), NCSD, USGS, and
27 Woodlands. The integration of GSE data is accomplished by using computer software to
28 interpolate between measurements and calculate GWI within the principal production aquifer
29 assuming an unconfined aquifer and a specific yield of 11.7 percent. Limited measurements of
30 GSE were available for the years 1982, 1983, 1984, 1994 and 1997, precluding a reliable
31 calculation of GWI for those years.

32 Groundwater Surface Elevation Measurements

33 Groundwater surface elevation data were obtained from SLO DPW, NCSD, USGS, and
34 Woodlands. SLO DPW measures GSE in monitoring wells during the spring (April) and the fall

F
I
N
A
L

To: Don Spagnolo
Re: Spring 2010 GWI
Date: July 15, 2010
Page: 2 of 5

1 (October) of each year. Woodlands and NCS D measures GSE in their monitoring wells
2 monthly. For the years 1975 to 1999, available representative GSE data were used to compute
3 GWI. For the years 2000 to 2010, only GSE data from the same 45 wells were used to compute
4 GWI.

5 The GSE data was reviewed in combination with well completion reports and historical
6 hydrographic records in order to exclude measurements that do not accurately represent static
7 water levels within the principal production aquifer. Wells that do not access the principal
8 production aquifer or were otherwise determined to not accurately represent static water levels
9 within the aquifer were not included in analysis.

10 **Groundwater Surface Interpolation**

11 The individual GSE measurements from each year were used to produce a GSE field by
12 interpolation using the inverse distance weighting (IDW) method.

13 **Groundwater Index**

14 The value of the groundwater index was computed for the area defined in Phase III of the
15 trial. The GWI was computed by subtracting both the mean sea level surface (elevation equals
16 zero) and the volume of bedrock above sea level from the hypothetical saturated volume. The
17 bedrock surface elevation is based on Figure 11: Base of Potential Water-Bearing Sediments,
18 presented in the report, Water Resources of the Arroyo Grande - Nipomo Mesa Area (DWR
19 2002). The bedrock surface elevation was preliminarily verified by reviewing driller reports
20 obtained from DWR. The saturated volume above sea level and bedrock was multiplied by a
21 specific yield of 11.7% to compute the GWI. The specific yield is based on the average weighted
22 specific yield measurement made at wells within the Nipomo Mesa Hydrologic Sub-Area (DWR
23 2002, pg. 86).

24 **Key Well Index**

25 The NMMA Technical Group selected the data from eight inland key wells to represent
26 the whole of the NMMA. The Key Well Index was calculated annually using Spring GSE
27 measurements from 1975 to 2008. The Key Wells were selected to represent various portions of
28 the groundwater basin within the NMMA. In selecting the eight key wells, the following
29 criteria were applied so that the wells generally represent the NMMA as a whole:

- 30 (1) The wells are geographically distributed,
- 31 (2) No single well overly influences the Key Well Index.

32 The first criterion was met in the selection of the wells, such that no well represented a
33 disproportionate area. To meet the second criterion, groundwater elevations from each well
34 were normalized so that any well where elevations were on the average higher or lower than
35 the other wells did not overly influence the magnitude of the Key Well Index. This

F
I
N
A
L

To: Don Spagnolo
Re: Spring 2010 GWI
Date: July 15, 2010
Page: 3 of 5

1 normalization was accomplished by dividing each spring groundwater elevation measurement
2 by the sum of all the Spring GSE data for that well.

3 The Key Well Index was defined for each year as the average of the normalized spring
4 groundwater data from each well. The lowest value of the Key Well Index could be considered
5 the "historical low" within the NMMA.

6

7 **REFERENCES**

8 Department of Water Resources (DWR). 2002. Water Resources of the Arroyo Grande -
9 Nipomo Mesa Area, Southern District Report.

F
I
N
A
L

To: Don Spagnolo
 Re: Spring 2010 GWI
 Date: July 15, 2010
 Page: 4 of 5

Table 1

Spring and Fall
 Groundwater Index
 (GWI)


Year	Rainfall (Inches)	Spring GWI (Acre-Feet)	Number of Wells	Fall GWI (Acre-Feet)	Number of Wells	Spring to Fall Difference (Acre-Feet)
1975	17.29	99,000	54	91,000	54	8,000
1976	13.45	82,000	45	76,000	65	6,000
1977	10.23	64,000	59	54,000	63	10,000
1978	30.66	84,000	62	---	35	---
1979	15.80	72,000	57	77,000	63	(5,000)
1980	16.57	88,000	55	89,000	46	(1,000)
1981	13.39	97,000	46	75,000	47	22,000
1982	18.58	123,000	42	---	31	---
1983	33.21	---	35	95,000	42	---
1984	11.22	---	14	76,000	37	---
1985	12.20	106,000	37	82,000	41	24,000
1986	16.85	98,000	51	67,000	51	31,000
1987	11.29	83,000	48	71,000	52	12,000
1988	12.66	80,000	51	66,000	49	14,000
1989	12.22	59,000	47	47,000	57	12,000
1990	7.12	62,000	55	49,000	53	13,000
1991	13.06	62,000	52	55,000	54	7,000
1992	15.66	61,000	52	35,000	48	26,000
1993	20.17	72,000	54	52,000	61	20,000
1994	12.15	60,000	54	---	36	---
1995	25.47	87,000	35	74,000	52	25,000
1996	16.54	76,000	45	62,000	57	14,000
1997	20.50	---	20	91,000	48	---
1998	33.67	105,000	41	93,000	44	12,000
1999	12.98	106,000	56	88,000	49	18,000
2000	14.47	108,000	44	84,000	41	24,000
2001	18.78	118,000	43	85,000	35	33,000
2002	8.86	96,000	29	79,000	41	17,000
2003	11.39	94,000	37	66,000	42	28,000
2004	12.57	89,000	42	81,000	35	8,000
2005	22.23	98,000	38	79,000	39	19,000
2006	20.83	107,000	44	78,000	41	29,000
2007	6.96	93,000	44	66,000	42	27,000
2008	15.18	83,000	43	65,000	42	18,000
2009	10.31	76,000	44	65,000	43	11,000
2010	17.05	80,000	45			

---: insufficient for evaluation

F
I
N
A
L

To: Don Spagnolo
 Re: Spring 2010 GWI
 Date: July 15, 2010
 Page: 5 of 5

Figure 1



F
I
N
A
L