TO:

**BOARD OF DIRECTORS** 

FROM:

MICHAEL S. LEBRUN MYC

INTERIM GENERAL MANAGER

DATE:

DECEMBER 9, 2011

AGENDA ITEM E-1

**DECEMBER 14, 2011** 

# RECEIVE GROUNDWATER INDEX PRESENTATION BY BRAD NEWTON, Ph.D, PG FROM WAGNER& BONSIGNORE CONSULTING CIVIL ENGINEERS

## ITEM

Presentation of the fall groundwater index for the Nipomo Mesa area. [Receive Report]

#### BACKGROUND

Brad Newton of Wagner & Bonsignore is scheduled to summarize the attached report. The report is an independent product of Wagner & Bonsignore and is not reviewed or recognized by the Nipomo Mesa Management Area Technical group.

## **FISCAL IMPACT**

Preparation of this report is included in the FY 2011-12 Budget.

### RECOMMENDATION

Staff recommends that the Board receive the Report and give direction to staff.

## **ATTACHMENTS**

Fall 2011 Groundwater Index

t:\board matters\board meetings\board letter\2011\111214 ground water index.docx

# TECHNICAL MEMORANDUM

2

7

8

9

10

11

12

13

14

15

16 17

18

19

1

3 TO:

NCSD Board of Directors

4 FROM:

Brad Newton, Ph.D., P.G.; Jesse Herbert

5 RE:

Fall 2011 Groundwater Index

6 DATE:

December 07, 2011

## INTRODUCTION

Groundwater surface elevations (GSE) underlying the Nipomo Mesa are regularly measured at many places (wells) across the mesa. The Fall 2011 Groundwater Index (GWI) has been computed and presented herein along with historical GWI from 1975 to present based on these groundwater surface elevation measurements collected during spring and fall across the Nipomo Mesa. Limited measurements of GSE were available for the years 1982, 1983, 1984, 1994 and 1997, thus precluding a reliable calculation of GWI for those years.

Ground elevation surveys for the key wells were conducted in preparation of the 1st Annual Report - Calendar Year 2008 for the Nipomo Mesa Management Area (NMMA). These updated reference points were not incorporated into the GWI to preserve consistency in the historical calculations and presentations.

The NMMA Technical Group has not reviewed this technical memorandum, its findings, or any presentation of this evaluation.

20 21

22

23

24

## RESULTS

Fall 2011 GWI is 81,000 acre-feet (AF), which is 14,000 AF greater than the Fall 2010 GWI (Table 1, Figure 1). The Key Well Index from NMMA 3rd Annual Report - Calendar Year 2010 generally follows the same historical trends as the GWI (Figure 1).

25 26

27

28

29

30

31

32

33

#### METHODOLOGY

The calculation of spring and fall GWI are based on GSE measurements regularly made by San Luis Obispo County Department of Public Works (SLO DPW), NCSD, USGS, and Woodlands. The integration of GSE data is accomplished by using computer software to interpolate between measurements and calculate GWI within the principal production aquifer assuming an unconfined aquifer and a specific yield of 11.7 percent. Limited measurements of GSE were available for the years 1982, 1983, 1984, 1994 and 1997, precluding a reliable calculation of GWI for those years.

s:\client files\ncsd (6001)\tasks\general consultation - 6001.2\activities\tm23 gwi fall 2011\deliverable\20111207 fall 2011 gwi.doc

RE: Fall 2011 GWI DATE: December 07, 2011

Page 2 of 6

### Groundwater Surface Elevation Measurements

Groundwater surface elevation data were obtained from SLO DPW, NCSD, USGS, and Woodlands. SLO DPW measures GSE in monitoring wells during the spring (April) and the fall (October) of each year. Woodlands and NCSD measures GSE in their monitoring wells monthly. For the years 1975 to 1999, available representative GSE data were used to compute GWI. For the years 2000 to 2011, only GSE data from the same 45 wells were used to compute GWI.

The GSE data was reviewed in combination with well completion reports and historical hydrographic records in order to exclude measurements that do not accurately represent static water levels within the principal production aquifer. Wells that do not access the principal production aquifer or were otherwise determined to not accurately represent static water levels within the aquifer were not included in analysis.

# Groundwater Surface Interpolation

The individual GSE measurements from each year were used to produce a GSE field by interpolation using the inverse distance weighting (IDW) method.

## Groundwater Index

The GWI is defined as the saturated volume above sea level and bedrock multiplied by the specific yield of 11.7 percent. The value of the groundwater index was computed for the area defined in Phase III of the trial. The base of the saturated volume is mean sea level surface (elevation equals zero) or the bedrock above sea level, whichever is higher. The bedrock surface elevation is based on Figure 11: Base of Potential Water-Bearing Sediments, presented in the report, Water Resources of the Arroyo Grande – Nipomo Mesa Area (DWR 2002). The bedrock surface elevation was preliminarily verified by reviewing driller reports obtained from DWR (Figure 2). The specific yield is based on the average weighted specific yield measurement made at wells within the Nipomo Mesa Hydrologic Sub-Area (DWR 2002, pg. 86).

# Key Well Index

The NMMA Technical Group selected the data from eight inland key wells to represent the whole of the NMMA. The Key Well Index was calculated annually using spring GSE measurements from 1975 to 2009. The key wells were selected to represent various portions of the groundwater basin within the NMMA. In selecting the eight key wells, the following criteria were applied so that the wells generally represent the NMMA as a whole:

- (1) The wells are geographically distributed,
- (2) No single well overly influences the Key Well Index.

The first criterion was met in the selection of the wells, such that no well represented a disproportionate area. To meet the second criterion, groundwater elevations from each well were normalized so that any well where elevations were on the average higher or lower than

R

A

F

T

|    | RE: Fall 2011 GWI   |
|----|---|
|    | DATE: December 07, 2011   |
|    | Page 3 of 6   |
| 1  | the other wells did not overly influence the magnitude of the Key Well Index. This          |
| 2  | normalization was accomplished by dividing each spring groundwater elevation measurement    |
| 3  | by the sum of all the Spring GSE data for that well.  |
| 4  | The Key Well Index was defined for each year as the average of the normalized spring        |
| 5  | groundwater data from each well. The lowest value of the Key Well Index could be considered |
| 6  | the "historical low" within the NMMA.   |
| 7  |   |
| 8  | REFERENCES  |
| 9  | Department of Water Resources (DWR). 2002. Water Resources of the Arroyo Grande - Nipomo    |
| 10 | Mesa Area, Southern District Report.  |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |

TO:

NCSD Board of Directors

RE: Fall 2011 GWI DATE: December 07, 2011

Page 4 of 6

# Spring and Fall Groundwater Index (GWI)

| Year | Rainfall<br>(inches) | Spring GWI<br>(Acre-Feet) | Number<br>of Wells | Fall GWI<br>(Acre-Feet) | Number<br>of Wells | Spring to Fall<br>Difference<br>(Acre-Feet) |
|------|----------------------|---------------------------|--------------------|-------------------------|--------------------|---|
| 1975 | 17.29                | 99,000                    | 54                 | 91,000                  | 54                 | 8,000                                       |
| 1976 | 13.45                | 82,000                    | 45                 | 76,000                  | 65                 | 6,000                                       |
| 1977 | 10.23                | 64,000                    | 59                 | 54,000                  | 63                 | 10,000                                      |
| 1978 | 30.66                | 84,000                    | 62                 | 2-4                     | 35                 | -   |
| 1979 | 15.80                | 72,000                    | 57                 | 77,000                  | 63                 | (5,000)                                     |
| 1980 | 16.57                | 88,000                    | 55                 | 89,000                  | 46                 | (1,000)                                     |
| 1981 | 13.39                | 97,000                    | 46                 | 75,000                  | 47                 | 22,000                                      |
| 1982 | 18.58                | 123,000                   | 42                 |                         | 31                 | -   |
| 1983 | 33.21                |                           | 35                 | 95,000                  | 42                 |   |
| 1984 | 11.22                |                           | 14                 | 76,000                  | 37                 | _   |
| 1985 | 12.20                | 106,000                   | 37                 | 82,000                  | 41                 | 24,000                                      |
| 1986 | 16.85                | 98,000                    | 51                 | 67,000                  | 51                 | 31,000                                      |
| 1987 | 11.29                | 83,000                    | 48                 | 71,000                  | 52                 | 12,000                                      |
| 1988 | 12.66                | 80,000                    | 51                 | 66,000                  | 49                 | 14,000                                      |
| 1989 | 12.22                | 59,000                    | 47                 | 47,000                  | 57                 | 12,000                                      |
| 1990 | 7.12                 | 62,000                    | 55                 | 49,000                  | 53                 | 13,000                                      |
| 1991 | 13.06                | 62,000                    | 52                 | 55,000                  | 54                 | 7,000                                       |
| 1992 | 15.66                | 61,000                    | 52                 | 35,000                  | 48                 | 26,000                                      |
| 1993 | 20.17                | 72,000                    | 54                 | 52,000                  | 61                 | 20,000                                      |
| 1994 | 12.15                | 60,000                    | 54                 |                         | 36                 |   |
| 1995 | 25.47                | 87,000                    | 35                 | 74,000                  | 52                 | 25,000                                      |
| 1996 | 16.54                | 76,000                    | 45                 | 62,000                  | 57                 | 14,000                                      |
| 1997 | 20.50                |                           | 20                 | 91,000                  | 48                 | -   |
| 1998 | 33.67                | 105,000                   | 41                 | 93,000                  | 44                 | 12,000                                      |
| 1999 | 12.98                | 106,000                   | 56                 | 88,000                  | 49                 | 18,000                                      |
| 2000 | 14.47                | 108,000                   | 44                 | 84,000                  | 41                 | 24,000                                      |
| 2001 | 18.78                | 118,000                   | 43                 | 85,000                  | 35                 | 33,000                                      |
| 2002 | 8.86                 | 96,000                    | 29                 | 79,000                  | 41                 | 17,000                                      |
| 2003 | 11.39                | 94,000                    | 37                 | 66,000                  | 42                 | 28,000                                      |
| 2004 | 12.57                | 89,000                    | 42                 | 81,000                  | 35                 | 8,000                                       |
| 2005 | 22.23                | 98,000                    | 38                 | 79,000                  | 39                 | 19,000                                      |
| 2006 | 20.83                | 107,000                   | 44                 | 78,000                  | 41                 | 29,000                                      |
| 2007 | 6.96                 | 93,000                    | 44                 | 66,000                  | 42                 | 27,000                                      |
| 2008 | 15.18                | 83,000                    | 43                 | 65,000                  | 42                 | 18,000                                      |
| 2009 | 10.31                | 76,000                    | 44                 | 65,000                  | 43                 | 11,000                                      |
| 2010 | 17.05                | 80,000                    | 45                 | 67,000                  | 42                 | 13,000                                      |
| 2011 | 29.19*               | 87,000                    | 43                 | 81,000                  | 43                 | 6,000                                       |

<sup>-:</sup> Insufficient for evaluation

<sup>\*:</sup> Preliminary value

Table 1: Groundwater Index computed from Spring 1975 to Fall 2011.

RE: Fall 2011 GWI DATE: December 07, 2011

Page 5 of 6

1 2




Figure 1: Groundwater Index from Spring 1975 to Fall 2011 and the Key Well Index computed from Spring 1975 to Spring 2010.

RE: Fall 2011 GWI DATE: December 07, 2011

Page 6 of 6

1 2

